

Quasiperiodic infinite words : multi-scale case and dynamical properties

Thierry Monteil* and Solomon Marcus†

Abstract

An infinite word x is said to be quasiperiodic if there exists a finite word q such that x is covered by occurrences of q (such a q is called a quasiperiod of x). Using the notion of derivation, we show that this definition is not sufficient to imply any symmetry in an infinite word. Therefore we introduce multi-scale quasiperiodic words, i.e. quasiperiodic words that admit an infinite number of quasiperiods. Such words are uniformly recurrent, this allows us to study the subshift they generate. We prove that multi-scale quasiperiodic subshifts are uniquely ergodic and have zero topological entropy as well as zero Kolmogorov complexity. Sturmian subshifts are shown to be multi-scale quasiperiodic.

Keywords: symbolic dynamics, subshift, quasiperiodicity, unique ergodicity, topological entropy, complexity, substitution, Rauzy graph, sturmian subshift.

AMS Classification: 68R15, 37B10, 37A35, 37A05.

*Institut de Mathématiques de Luminy, CNRS UMR 6206, Case 907, 163 Avenue de Luminy, 13288 Marseille cedex 09, France – monteil@iml.univ-mrs.fr

†Romanian Academy, Mathematics, Calea Victoriei 125, Bucharest, Romania – Solomon.Marcus@imar.ro

1 Introduction and notations

1.1 Background: quasiperiodicity and symmetry

The general concept of symmetry is central in the study of infinite words, and many notions are devoted to measure it, like word complexity, recurrence, Kolmogorov's complexity, entropy, ... Those notions are more or less linked and in any case the most symmetric words are the periodic ones.

In [Mar1], the second author introduced a new notion to describe symmetric words: quasiperiodicity. This notion is an extension of the notion of quasiperiodicity for finite words that was introduced in [ApoEhr] and has also roots in different contexts like musicology [CrIIra] and molecular biology [KMGL] [MilJur].

Let A be a finite set (called *alphabet*). Let $x = x_0x_1x_2\dots$ in $A^{\mathbb{N}}$ be an infinite word. We will say that x is *quasiperiodic* if there exists a finite word q in A^* such that x is covered by the occurrences of q (A^* denotes the set of finite words on the alphabet A). Such a q is called a *quasiperiod* of x ; it is a prefix of x .

More precisely, let $(i_n)_{n \in \mathbb{N}}$ be the increasing sequence whose image is the set $\{i \in \mathbb{N} \mid x_{i \rightarrow i+l(q)-1} = q\}$, where $l(q)$ denotes the length of q , and $x_{i \rightarrow j}$ denotes $x_i x_{i+1} x_{i+2} \dots x_j$ (for $0 \leq i \leq j$). The word q is a quasiperiod of x whenever (i_n) is a well defined infinite sequence such that $i_0 = 0$ and $i_{n+1} - i_n \leq l(q)$ for any integer n .

For example, the infinite word $x = ababaabaabaababababaabababaabaabaabaabaaba\dots$ is quasiperiodic with aba as a quasiperiod and the infinite quasiperiodic words that admit aba as a quasiperiod are precisely the words over the alphabet $\{a, b\}$ beginning with the word ab and in which the words aaa and bb do not appear.

The second author was then looking for relations with other notions of symmetry. In [Mar1] and [LevRic1], it is shown that there is no such relation. Let us introduce a basic tool that will unify those results and allow us to understand this lack of fitting between quasiperiodicity and other notions: the derivation.

1.2 The derivation: a change of scale

1.2.1 Derivation

Let $x = x_0x_1x_2\dots$ be a quasiperiodic word, and q be a quasiperiod of x . Let $(i_n)_{n \in \mathbb{N}}$ be the sequence as defined before. We can define $\frac{\partial x}{\partial q}$ to be the infinite word on the alphabet $\{0, \dots, l(q) - 1\}$ whose k^{th} letter is $l(q) - i_{k+1} + i_k$ ($k \geq 0$).

In other terms, the k^{th} letter of $\frac{\partial x}{\partial q}$ is the length of the overlap between the k^{th} and the $(k+1)^{\text{th}}$ occurrence of q in x .

For example, if $x = ababaabaabaababababaababababaabaabaabaabaaba\dots$ is quasiperiodic with aba as a quasiperiod, then $\frac{\partial x}{\partial aba} = 100011101100010\dots$

In some sense, the $\frac{\partial}{\partial q}$ operator removes exactly the information contained in q since the knowledge of $\frac{\partial x}{\partial q}$ and q is sufficient to reconstruct x .

For example, if $x = 01121010201\dots$ and $w = abcaa$,
then $\int_w x = abcaaaabcaaabcaabcaabcaaaaabcaabcaaaaabcaabcaaaaabcaabcaa....$

1.2.3 Quasiperiodicity is spread everywhere

To construct a non recurrent quasiperiodic word, just take a non recurrent word x on the alphabet $\{0, 1\}$ and consider $\int_{aba} x$. Do the same to construct a uniformly recurrent quasiperiodic word.

Complexity For any infinite word x and any integer n , $L_n(x)$ denotes the finite words of length n that occur in x and we define $L(x) \stackrel{\text{def}}{=} \bigcup_{n \in \mathbb{N}} L_n(x)$.

We can define an equivalence relation on the set of infinite words through the asymptotic behaviour of their complexity function: two words x and y are said to be *complexity equivalent* if there exists a positive integer K such that for all $n \geq 1$, $p_n(x) \leq Kp_{Kn}(y)$ and $p_n(y) \leq Kp_{Kn}(x)$. Hence, bounded, linear, quadratic, polynomial or exponential growths are preserved under complexity equivalence.

Proof: Let x be an infinite word on an alphabet A . There is no restriction to suppose that $A = \{0, \dots, k\}$. Let $y \stackrel{\text{def}}{=} \int_{a^k b a^k} x$. We check that y is a quasiperiodic word in the class of complexity equivalence of x . $\square$

3

In [LevRic1], Florence Levé and Gwénaél Richomme proved that there exists sturmian words that are not quasiperiodic.

Entropy is the exponent of the complexity function in the vocabulary of topological dynamical systems (see section 3).

Frequencies If u and v are finite words, let $\#(u, v)$ denotes the number of occurrences of u in v . An infinite word x is said to have *frequencies* if for any finite word w , $\frac{1}{n}\#(w, x_{0 \rightarrow n-1})$ admits a limit when n tends to infinity.

By integration, there exists quasiperiodic words that do not have frequencies: if x is an infinite word such that 0 does not appear with frequencies, then $\int_{aba} x$ is a quasiperiodic word such that bab does not appear with frequencies.

Unique ergodicity is a strong analogue of having frequencies in the vocabulary of topological dynamical systems (see section 4).

Hence the notion of quasiperiodicity does not insert well among other notions of symmetry.

1.3 Multi-scale quasiperiodic words

We have to notice that all classical notions of symmetry are invariant under such a renormalization procedure (derivation corresponds to induction if we are studying dynamical properties like entropy [Abr]). Therefore, if we want that a notion says something about symmetry, we should ensure that it is stable under such a change of scale. As we saw with the derivation procedure, the existence of a quasiperiod q in a word x just imposes rigidity at the scale around $l(q)$ but does not impose anything at larger scales.

This leads to the following definition : an infinite word is said to be *multi-scale quasiperiodic* if the set $Q(x)$ of its quasiperiods is infinite.

The easiest non-periodic multi-scale quasiperiodic words we can construct, are the fixed points for some particular integration operators. For example, the fixed point of $\int_{010}$ is multi-scale quasiperiodic, it is known as the Fibonacci word (a precise description of the quasiperiods of this word can be found in [LevRic1]). There exists much wilder multi-scale quasiperiodic words (see Theorem 4 in section 3).

In the next section we will prove that multi-scale quasiperiodic words are uniformly recurrent. This will allow us to study the subshift generated by them (section 2). Concerning the complexity, we will prove that multi-scale quasiperiodic subshifts have zero topological entropy as well as zero Kolmogorov complexity (section 3). We will also prove that sturmian subshifts are multi-scale quasiperiodic (section 5). Concerning frequencies, we will prove that multi-scale quasiperiodic subshifts are uniquely ergodic (section 4).

2 Uniform recurrence and minimality

An infinite word $x \in A^{\mathbb{N}}$ is said to be *uniformly recurrent* if any finite word $u \in L(x)$ occurs infinitely many times in x and the gap between two consecutive occurrences of u in x is bounded, equivalently

$$\forall u \in L(x) \quad \exists n \geq 1 \quad \forall v \in L_n(x) \quad \#(u, v) \geq 1$$

Theorem 2. *Any multi-scale quasiperiodic word x is uniformly recurrent.*

Proof: Let u be a finite word that occurs in x . Since every quasiperiod of x is a prefix of x , one of them must contain an occurrence of u (they have unbounded length). Let q be such a quasiperiod. Any word in $L_{2l(q)}$ contains at least an occurrence of q and therefore at least an occurrence of u . $\square$

This property of multi-scale quasiperiodic words allows us to deal with the dynamical system generated by a multi-scale quasiperiodic word as follows:

We endow A with the discrete topology and $A^{\mathbb{N}}$ with the product topology. This makes $A^{\mathbb{N}}$ a metrisable compact space.

We note

$$S \stackrel{\text{def}}{=} \left(\begin{array}{ccc} A^{\mathbb{N}} & \longrightarrow & A^{\mathbb{N}} \\ x = x_0 x_1 \dots x_n \dots & \longmapsto & x_1 x_2 \dots x_{n+1} \dots \end{array} \right)$$

for the shift. It is a continuous map.

If x is a multi-scale quasiperiodic word, we define

$$X \stackrel{\text{def}}{=} \overline{\{S^k(x) \mid k \in \mathbb{N}\}}$$

and we still note S for the restriction of S to X , making (X, S) a topological dynamical system.

Theorem 2 is equivalent to say that (X, S) is a *minimal subshift* i.e. X is a nonempty closed subset of $A^{\mathbb{N}}$ stable under S and that is minimal for those properties.

A minimal subshift generated by a multi-scale quasiperiodic word is called a *multi-scale quasiperiodic subshift*.

If u is a finite word, we define the *cylinder*

$$[u] \stackrel{\text{def}}{=} \{x \in X \mid (\forall i \leq n-1)(x_i = u_i)\} \quad (u = u_0 \dots u_{n-1} \in A^n \quad (n \in \mathbb{N}))$$

We can notice that the derivation of a multiscaled quasiperiodic word over the quasiperiod q corresponds to the induction of the subshift (X, S) on the cylinder $[q]$.

We extend the notion of language and complexity to minimal subshifts:

$$L_n(X) \stackrel{\text{def}}{=} \{u \in A^n \mid [u] \neq \emptyset\} \quad \text{and} \quad p_n(X) = \text{card}(L_n(X)) \quad (n \in \mathbb{N})$$

$$L(X) \stackrel{\text{def}}{=} \bigcup_{n \in \mathbb{N}} L_n(X)$$

If y is in X , we have $L(y) = L(X)$ and $p.(y) = p.(X)$.

3 Complexity and topological entropy

3.1 Word complexity

Theorem 3. *Let x be a multi-scale quasiperiodic word. Then*

$$\liminf_{n \rightarrow \infty} \frac{p_n(x)}{n^2} \leq 1 < \infty$$

Proof: Let $q \in Q(x)$ and $u \in L_{l(q)}(x)$. Since q is a quasiperiod of x , u is a subword of some $v.q$ where v is nonempty prefix of q . u is determined by the choice of v and his position in $v.q$. There are $l(q)$ prefixes of q and for such a prefix v , there are $l(v) \leq l(q)$ available positions for u (we do not count q several times). Finally, there are only $l(q)^2$ possibilities for u and $p_{l(q)}(x) \leq l(q)^2$. The result follows since $\{l(q) \mid q \in Q(x)\}$ is not bounded. $\square$

Theorem 4. *For each positive function $f : \mathbb{N} \rightarrow \mathbb{R}_+^*$ that converges to zero, there exists a multi-scale quasiperiodic subshift (X, S) such that $\frac{1}{n} \log(p_n(X)) \geq f(n)$ for infinitely many n . In particular, we can ask to $p_n(X)$ to grow faster than any polynomial on a subsequence.*

Proof: Let $\varphi : \mathbb{N} \rightarrow \mathbb{N}^*$ be an increasing sequence such that $2nf(2n\varphi(n)) \leq 1$ for any n . Let $A = \{0, 1\}$ be the alphabet. For $n \geq 1$, let w_n be a finite word on A such that

- w_n begins and ends with the letter 0,
- every word w of length $2n$ such that $\#(0, w) = \#(1, w) = n$ appears as a subword of w_n .

Now, let us define by induction a sequence of finite words over A :

- $u_0 = 010$
- $u_{n+1} = \int_{u_n} w_{\varphi(l(u_n))}$

which is well defined since for any n , u_n begins and ends with the letter 0.

Since $u_n = \int_{u_n} 0$, u_n is a prefix of u_{n+1} : let x be the unique infinite word over A such that u_n is a prefix of x for any n .

If $k \leq l$, u_l is covered by occurrences of u_k , so all the u_n are quasiperiods of x and x is multi-scale quasiperiodic. Let (X, S) denote the associated minimal subshift

Let n be a positive integer. u_{n+1} and therefore x contain an occurrence of $\int_{u_n} w$, where w is any word of length $2\varphi(l(u_n))$ such that $\#(0, w) = \#(1, w) = \varphi(l(u_n))$. There are at least $2^{\varphi(l(u_n))}$ such different words and each of them has length $\varphi(l(u_n))(l(u_n) + (l(u_n) - 1))$.

Hence,

$$p_{2\varphi(l(u_n))l(u_n)}(X) \geq p_{\varphi(l(u_n))(l(u_n)+(l(u_n)-1))}(X) \geq 2^{\varphi(l(u_n))}$$

So,

$$\frac{1}{2\varphi(l(u_n))l(u_n)} \log(p_{2\varphi(l(u_n))l(u_n)}(X)) \geq \frac{\varphi(l(u_n))}{2\varphi(l(u_n))l(u_n)} = \frac{1}{2l(u_n)} \geq f(2\varphi(l(u_n))l(u_n))$$

$\square$

We can notice that for the infinite words constructed here, the scales controlled by the quasiperiods are very sparse.

3.2 Topological entropy

If (X, S) is a minimal subshift, then the limit

$$h_{top}(X) \stackrel{\text{def}}{=} \lim_{n \rightarrow \infty} \frac{1}{n} \log(p_n(X))$$

exists and is named the *topological entropy* of X .

Corollary 1. *Any multi-scale quasiperiodic subshift has zero topological entropy.*

3.3 Kolmogorov complexity

Let U be a fixed universal Turing machine and for each finite word $u \in A^*$, let $K_U(u)$ denotes the Kolmogorov complexity associated to u , i.e. the length of the shortest binary word p such that $U(p) = u$ (see [Bru]). For a minimal subshift (X, S) , we can define

$$K(X) \stackrel{\text{def}}{=} \sup_{x \in X} \limsup_{n \rightarrow \infty} \frac{K_U(x_{1 \rightarrow n})}{n}$$

(this number is independant of the choice of U since if U' is another universal Turing machine, there is a constant C such that for any finite word u , $K_U(u) - C \leq K_{U'}(u) \leq K_U(u) + C$).

Corollary 2. *For any multi-scale quasiperiodic subshift (X, S) , $K(X) = 0$.*

Indeed, [Bru] Theorem 3.1 asserts that $K(X) \leq h_{top}(X)$.

But we can also give a more direct argument : if q is a fixed quasiperiod of a multi-scale quasiperiodic subshift (X, S) and if x is in X , then $\limsup_{n \rightarrow \infty} \frac{K_U(x_{1 \rightarrow n})}{n} \leq \frac{4 \log(l(q))}{l(q)}$. To prove this, it suffices to remark that the integration algorithm can be coded in $O(1)$.

Then, if n is bigger than $4l(q)$, there exists three finite words u , w and v such that $x_{1 \rightarrow n} = u.w.v$, the lengths of u and v are uniformly bounded by $l(q)$, and w begins and ends with q (and can therefore be derivated).

The remaining problem is the control of the length of $\partial w / \partial q$: if the occurrences of q overlap each other deeply, $\partial w / \partial q$ can be rather long. To solve this, we decide to replace recursively an occurrence of $n_1.n_2$ in $\partial w / \partial q$ by $n_1 + n_2$ if n_1 and n_2 are smaller than $n/2$. This operation consists in omitting some useless occurrences of q in $x_{1 \rightarrow n}$. Then at least half of the numbers appearing in the new form of $\partial w / \partial q$ are bigger than $l(q)/2$, so the length of this new $\partial w / \partial q$ will be less than $4n/l(q)$.

Since the coding of each letter of $\partial w / \partial q$ costs $\log(l(q))$, we have a total cost less than a constant (to code the integration algorithm, q , u and v) plus $\log(l(q))4n/l(q)$ (to code the new form of $\partial w / \partial q$).

4 Unique ergodicity and frequencies

For a multi-scale quasiperiodic subshift (X, S) , we will now study the set $\mathcal{M}(X, S)$ of Borel probability measures on X that are invariant under S . This set can be identified with a nonempty compact convex subset of $C^0(X, \mathbb{R})'$ endowed with the weak-star topology.

A S -invariant measure $\mu \in \mathcal{M}(X, S)$ is said to be *ergodic* if the only Borel sets $A \subset X$ such that $S^{-1}(A) = A$ have measure $\mu(A) = 0$ or 1 . Such measures are the extremal points of $\mathcal{M}(X, S)$ and satisfies Birkhoff's theorem :

$$\forall f \in L^1(X, \mathbb{R}) \quad \frac{1}{n} \sum_{k=0}^{n-1} f \circ S^k \xrightarrow[n \rightarrow \infty]{\mu\text{-a.e.}} \int_X f d\mu$$

A minimal subshift is said to be *uniquely ergodic* if $\text{card}(\mathcal{M}(X, S)) = 1$.

One interest of such a situation is that, the unique invariant measure μ is ergodic, moreover the convergence in Birkhoff's theorem is uniform for continuous functions.

Theorem 5. *Any multi-scale quasiperiodic subshift (X, S) is uniquely ergodic.*

Proof:

We will first construct a S -invariant probability measure on X and then prove that it is the only one.

Step 1: We construct a candidate to be the unique measure. For this, we will approximate X by periodic subshifts generated by the $q^\omega = qqq \dots$ for $q \in Q(X)$.

For q in $Q(X)$, let

$$\mu_q \stackrel{\text{def}}{=} \frac{1}{l(q)} \sum_{k=0}^{l(q)-1} \delta_{S^k(q^\omega)}$$

(δ stands for the one-point Dirac's measure).

μ_q is the only element of $\mathcal{M}(A^\mathbb{N}, S)$ that gives measure 1 to the periodic subshift generated by the periodic word q^ω . By compacity we can find an infinite subset $Q' \subset Q(X)$ such that

$$\mu_q \xrightarrow[l(q) \rightarrow \infty]{q \in Q'} \mu$$

for some μ in $\mathcal{M}(A^\mathbb{N}, S)$.

Note that if X is aperiodic the μ_q 's give measure 0 to X . However μ will give strictly positive measure to X (as we will see in Step 2) : it shouldn't be surprising since the characteristic function of X is not continuous.

Step 2: Let us show that $\mu(X) = 1$. Since X is closed, we have the following approximation by clopen sets:

$$X = \overline{X} = \bigcap_{n \geq 1} \bigcup_{u \in L_n(X)} [u].$$

Let $n \geq 1$ and let $q \in Q'$ such that $l(q) \geq n$. For $i \in \{0, \dots, l(q) - n\}$, we have $q_{i \rightarrow i+n-1}^\omega \in L_n(X)$ (as a subword of q). Hence $\mu_q(\bigcup_{u \in L_n(X)} [u]) \geq (l(q) - n + 1)/l(q)$.

Letting $l(q)$ tending to infinity, since the characteristic function of $\bigcup_{u \in L_n(X)} [u]$ is continuous, we have $\mu(\bigcup_{u \in L_n(X)} [u]) = 1$. By countable intersection (n is arbitrary), we have

$\mu(X) = 1$. Hence, we can still denote by μ for the restriction of μ to X .

Step 3: Let ν be an ergodic measure on X . We will show that $\nu = \mu$
By Birkhoff's theorem, there is x in X such that for $u \in L(X)$,

$$\nu([u]) = \lim_{n \rightarrow \infty} \frac{1}{n} \#(u, x_{0 \rightarrow n+l(u)-2}) = \lim_{n \rightarrow \infty} \frac{1}{n} \#(u, x_{0 \rightarrow n-1})$$

Let q in Q' such that $l(q) \geq l(u)$. We decompose x into blocks of length $2l(q)$: $x = B_0.B_1.B_2.B_3.B_4 \dots$ with $B_i = x_{2l(q)i \rightarrow 2l(q)(i+1)-1}$. Since each B_i is in $L_{2l(q)}(X)$, it contains at least one occurrence of q , hence $\#(u, B_0.B_1.B_2 \dots B_i) \geq (i+1)\#(u, q)$ for each i in $\mathbb{N}^*$. We have

$$\nu([u]) = \lim_{i \rightarrow \infty} \frac{1}{2l(q)i} \#(u, B_0.B_1.B_2 \dots B_i) \geq \lim_{i \rightarrow \infty} \frac{i+1}{2l(q)(i+1)} \#(u, q) = \frac{1}{2l(q)} \#(u, q)$$

Moreover, we can control the frequency of occurrences of u in q^ω by estimating the occurrences of u in q and bounding the number of occurrences of u that appear between two consecutive occurrences of q in q^ω :

$$\mu_q([u]) = \lim_{n \rightarrow \infty} \frac{1}{n} \#(u, q_{0 \rightarrow n+l(u)-2}^w) \leq \frac{1}{l(q)} (\#(u, q) + l(u)) = \frac{1}{l(q)} \#(u, q) + \frac{l(u)}{l(q)}$$

Therefore,

$$\mu_q([u]) \leq \frac{1}{l(q)} \#(u, q) + \frac{l(u)}{l(q)} \leq 2\nu([u]) + \frac{l(u)}{l(q)}$$

Letting $l(q)$ tend to infinity, we have $\mu([u]) \leq 2\nu([u])$. So, μ is absolutely continuous relatively to ν . It is well known that this implies $\mu = \nu$, but for sake of completeness, we include a short proof here. There exists a measurable function $f \in L^1(X, \mathbb{R}_+)$ such that for each borel set $A \subset X$, $\mu(A) = \int_A f d\nu$. Since μ is S -invariant, we have $\int_A f d\mu = \int_{S^{-1}A} f d\mu$ for each borel set $A \subset X$.

Let us show that f is constant almost everywhere. Assume by contradiction that the measure of set $A \stackrel{\text{def}}{=} \{x \in X / f(x) \geq \int_X f d\nu\}$ is in $]0, 1[$. Since ν is S -ergodic, A is not S -invariant, so $\nu(S^{-1}(A) \setminus A) = \nu(A \setminus S^{-1}(A)) > 0$.

Hence $\nu(A \setminus S^{-1}(A)) \int_X f d\nu \leq \int_{A \setminus S^{-1}(A)} f d\nu = \int_A f d\nu - \int_{A \cap S^{-1}(A)} f d\nu = \int_{S^{-1}(A)} f d\nu - \int_{A \cap S^{-1}(A)} f d\nu = \int_{S^{-1}(A) \setminus A} f d\nu < \nu(A \setminus S^{-1}(A)) \int_X f d\nu$ which is absurd.

Hence f is constant with value $\nu(X) = 1$, so $\mu = \nu$ and μ is the only S -invariant measure. $\square$

Corollary 3. *Let x be a multi-scale quasiperiodic word. Then each finite word u occuring in x has frequencies i.e. $\frac{1}{n} \#(u, x_{0 \rightarrow n-1})$ converges when $n \rightarrow \infty$.*

Proof: Let μ be the unique S -invariant measure for the associated subshift (X, S) . The characteristic function $\chi_{[u]}$ of $[u]$ is continuous, so we have a uniform and therefore a pointwise convergence in Birkhoff's theorem :

$$\frac{1}{n} \#(u, x_{0 \rightarrow n-1}) = \frac{1}{n} \sum_{k=0}^{n-1} \chi_{[u]}(S^k(x)) \xrightarrow{n \rightarrow \infty} \mu([u])$$

$\square$

We can remark that this proof has the same flavour as the the main result of [Bos] that asserts that every subshift with subaffine complexity has only a finite number of ergodic measures. In fact, Theorem 5 and Boshernitzan's result can both be deduced from a more general statement that involves the geometry of Rauzy graphs associated to a minimal subshift:

To each minimal subshift (X, S) we can associate a sequence $(G_n)_{n \geq 1}$ of oriented graphs as follows: the vertices of G_n is $L_n(X)$ and there is an edge from u to v if and only if there exists w in $L_{n+1}(X)$ such that u is a prefix of w and v is a suffix of w .

Those graphs are named the *Rauzy graphs* associated to (X, S) .

If $K \geq 1$, (X, S) is said to be *K-deconnectable* if there exists an extraction $\alpha \in \uparrow(\mathbb{N}^*, \mathbb{N}^*)$ and a constant $K' \geq 1$ such that for all $n \geq 1$ there exists a subset $D_{\alpha(n)} \subset L_{\alpha(n)}(X)$ of at most K vertices such that every path in $G_{\alpha(n)}(X) \setminus D_{\alpha(n)}$ is of length at most $K'\alpha(n)$ (in particular it do not contains any cycle). This means that, up to extraction, we can disconnect (in a specific way) the Rauzy graphs by removing at most K vertices.

Theorem 6 ([Mon]). *A K-deconnectable minimal subshift has at most K S-invariant ergodic measures.*

This result implies Boshernitzan's one, by taking for D_n the set of right special factors (i.e. the set of vertices having outgoing degree strictly greater than one). It also implies Theorem 5 since every multi-scale quasiperiodic subshift is 1-deconnectable. Indeed, if q is in $Q(X)$, $G_{l(q)}(X) \setminus \{q\}$ does not contain any path of length greater than $l(q)$.

5 Sturmian subshifts are multi-scale quasiperiodic

In [LevRic1], Florence Levé and Gwénaél Richomme proved that there exists sturmian words that are not quasiperiodic. In terms of complexity, sturmian words are the more symmetric words after periodic ones, so this result do not confirm that quasiperiodicity fits well with other notions of symmetry.

The dynamical point of view will allow us to solve the problem:

Theorem 7. *Sturmian subshifts are multi-scale quasiperiodic.*

Proof: Let (X, S) be a sturmian subshift. Since $p_1(X) = 2$, we can consider that X is defined on the alphabet $\{a, b\}$.

A word $u \in L_n X$ is said to be *left special* if au and bu are in $L_{n+1}(X)$. Since $p_{n+1}(X) - p_n(X) = 1$, there exists exactly one left special word l_n of length n ($n \geq 0$). A prefix of a left special word is still a left special word, so l_{n+1} begins with l_n : let us denotes by x the infinite word that begins by l_n for any integer n . Since $L(x) \subset L(X)$, x is in X (remember that X is closed). We will prove that x is multi-scale quasiperiodic.

The evolution of the Rauzy graphs of sturmian subshifts is described by Rauzy (see [ArnRau]): for infinitely many n (named “bursts”), the Rauzy graph $G_n(X)$ is eight shaped i.e. $G_n(X)$ is the union of two disjoint loops from l_n to l_n . By minimality, the minimal size of the two loops tends to infinity with n , in particular, it is positive for n big enough. Since the sum of the lengths of the two loops is equal to $p_n(X) = n + 1$, then each loop has size less or equal than n .

So, for infinitely many n , any path in $G_n(X)$ of length n starting from l_n has to meet l_n again i.e. l_n is a quasiperiod of x (x can be viewed as an infinite path in $G_n(X)$ starting from l_n). Therefore, (X, S) is a multi-scale quasiperiodic subshift.

□

Recently, Florence Levé and Gwénaél Richomme gave a precise description of the quasiperiodic sturmian words ([LevRic2]).

6 Conclusion

The gain of fitting with other symmetry classes obtained by considering multi-scale quasiperiodic subshifts instead of quasiperiodic words can be summarized as follow :




Figure 1: Quasiperiodic words among other words.




Figure 2: multi-scale quasiperiodic subshifts among other subshifts.

References

- [Abr] L.M. ABRAMOV, *The entropy of an induced transformation*, Doklady Akad. Nauk SSSR 128 (1959), p 647–650
- [ApoEhr] A. APOSTOLICO, A. EHRENFEUCHT, *Efficient detection of quasiperiodicities in strings*, Theoret. Comput. Sci. 119 (1993), no. 2, p 247–265.
- [ArnRau] P. ARNOUX, G. RAUZY, *Représentation géométrique de suites de complexité $2n + 1$* , Bull. Soc. Math. France 119 (1991), no. 2, p 199–215.
- [Bos] M. BOSHERNITZAN, *A unique ergodicity of minimal symbolic flows with linear block growth*, J. Analyse Math. 44 (1984/85), p 77–96.
- [Bru] A.A. BRUDNO, *Entropy and the complexity of the trajectories of a dynamic system*, Trudy Moskov. Mat. Obshch. 44 (1982), p 124–149.
- [CrIlRa] T. CRAWFORD, C. S. ILIOPOULOS, R. RAMAN, *String Matching Techniques for Musical Similarity and Melodic Recognition*, Computing in Musicology, vol. 11, p 73–100, 1998.
- [KMGL] S. KARLIN, M. MORRIS, G. GHANDOUR, M. LEUNG, *Efficient algorithms for molecular sequence analysis*, Proc. National. Acad. Sci U.S.A. 85 (1988), p 841–845.
- [LevRic1] F. LEVÉ, G. RICHOMME, *Quasiperiodic infinite words: some answers*, Bull. Eur. Assoc. Theor. Comput. Sci. EATCS No. 84, (2004), p 128–138.
- [LevRic2] F. LEVÉ, G. RICHOMME, *Private communication, april 2005*.
- [Mar1] S. MARCUS, *Quasiperiodic infinite words*, Bull. Eur. Assoc. Theor. Comput. Sci. EATCS No. 82, (2004), p 170–174.
- [Mar2] S. MARCUS, *Symmetry phenomena in infinite words, with biological, philosophical and aesthetic relevance*, <http://symmetry.hu/contents/contents-2003-2004.pdf>
- [MilJur] A. MILOSAVLJEVIC, J. JURKA, *Discovering simple DNA sequences by the algorithmic significance method*, Comput. Appl. Biosci., 9, (1993), p 407–411.
- [Mon] T. MONTEIL, *Estimating the number of ergodic measures of a minimal subshift knowing the geometry of its Rauzy graphs*, preprint.